


AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions and listings of claims in the application:

LISTING OF CLAIMS:

1. (currently amended): A method for treating obesity in a mammalian subject, which comprises ~~administration of an~~ administering to a mammalian subject in need of reduction of body weight an effective amount of a prostaglandin compound as shown by the following formula (I):




wherein L, M and N are hydrogen atom, hydroxy, halogen atom, lower alkyl, hydroxy(lower)alkyl, lower alkanoyloxy or oxo, wherein at least one of L and M is a group other than hydrogen, and the five-membered ring may have at least one double bond;

A is -CH₃, or -CH₂OH, -COCH₂OH, -COOH or a salt, ether, ester or amide thereof;

B is single bond, -CH₂-CH₂-, -CH=CH-, -C≡C-, -CH₂-CH₂-CH₂-, -CH=CH-CH₂-, -CH₂-CH=CH-, -C≡C-CH₂- or -CH₂-C≡C-;

Z is C=O



~~wherein R₄ and R₅ are hydrogen, hydroxy, halogen, lower alkyl, lower alkoxy or hydroxy(lower)alkyl, wherein R₄ and R₅ are not hydroxy and lower alkoxy at the same time;~~

R₁ is a saturated or unsaturated bivalent lower or medium aliphatic hydrocarbon residue, which is unsubstituted or substituted with halogen, alkyl, hydroxy, oxo, aryl selected from the group consisting of phenyl, tolyl and xylyl which is unsubstituted or substituted or heterocyclic group selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, furazanyl, pyranyl, pyridyl, pyridazinyl, pyrimidyl, pyrazinyl, 2-pyrrolinyl, pyrrolidinyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, piperidino, piperazinyl, morpholino, indolyl, benzothienyl, quinolyl, isoquinolyl, purinyl, quinazolinyl, carbazolyl, acridinyl, phenanthridinyl, benzimidazolyl, benzimidazolinyl, benzothiazolyl and phenothiazinyl which is unsubstituted or substituted, and at least one carbon atom in the aliphatic hydrocarbon is optionally substituted by oxygen, nitrogen or sulfur; and

R_a is a saturated or unsaturated lower or medium aliphatic hydrocarbon residue, which is unsubstituted or substituted with one or more substituents selected from the group consisting of halogen, oxo, hydroxy, lower alkoxy, lower alkanoyloxy, cyclo(lower)alkyl, cyclo(lower)alkyloxy, aryl, aryloxy, heterocyclic group and heterocyclic-oxy group; lower alkoxy; lower alkanoyloxy; cyclo(lower)alkyl; cyclo(lower)alkyloxy; aryl; aryloxy; heterocyclic group; or heterocyclic-oxy, ~~provided that Z is C=O and/or R_a is a saturated or unsaturated lower or medium aliphatic hydrocarbon residue which has a halogen substituent.~~

2. (canceled).
3. (currently amended): The method as described in Claim ~~4~~ 18, wherein said prostaglandin compound is a 16-mono or dihalogen-prostaglandin compound.
4. (currently amended): The method as described in Claim ~~4~~ 18, wherein said prostaglandin compound is a 13,14-dihydro-16-mono or dihalogen-prostaglandin compound.
5. (previously presented): The method as described in Claim 1, wherein said prostaglandin compound is a 13,14-dihydro-15-keto-16-mono or dihalogen-prostaglandin compound.
6. (currently amended): The method as described in Claim ~~4~~ 18, wherein said prostaglandin compound is a 13,14-dihydro-16-mono or difluoro-prostaglandin compound.
7. (previously presented): The method as described in Claim 1, wherein said prostaglandin compound is a 13,14-dihydro-15-keto-16-mono or difluoro-prostaglandin compound.
8. (currently amended): The method as described in Claim ~~4~~ 18, wherein said prostaglandin compound is a 13,14-dihydro-16-mono or dihalogen-prostaglandin E compound.
9. (previously presented): The method as described in Claim 1, wherein said prostaglandin compound is a 13,14-dihydro-15-keto-16-mono or dihalogen-prostaglandin E compound.
10. (currently amended): The method as described in Claim ~~4~~ 18, wherein said prostaglandin compound is a 13,14-dihydro-16,16-difluoro-prostaglandin E₁ compound.

11. (previously presented): The method as described in Claim 1, wherein said prostaglandin compound is a 13,14-dihydro-15-keto-16,16-difluoro-prostaglandin E₁ compound or 13,14-dihydro-15-keto-16,16-difluoro-18-methyl-prostaglandin E₁ compound.

12. (original): The method as described in Claim 1, which comprises systemic administration 1-4 times per day or continuous administration at the amount of 0.01-100 µg/kg per day.

13. (original): The method as described in Claim 12, wherein the administration is at the amount of 0.1-10 µg/kg per day.


14. (canceled).

15. (canceled).

16. (canceled).

17. (canceled).

18. (new): A method for treating obesity in a mammalian subject which comprises administering to a mammalian subject in need of reduction of body weight an effective amount of a prostaglandin compound as shown by the following general formula (I):




wherein L, M and N are hydrogen atom, hydroxy, halogen atom, lower alkyl, hydroxy(lower)alkyl, lower alkanoyloxy or oxo, wherein at least one of L and M is a group other than hydrogen, and the five-membered ring may have at least one double bond;

A is $-\text{CH}_3$, or $-\text{CH}_2\text{OH}$, $-\text{COCH}_2\text{OH}$, $-\text{COOH}$ or a functional derivative salt, ether, ester or amide thereof;

B is single bond, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-$, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-\text{CH}_2-$, $-\text{CH}_2-\text{CH}=\text{CH}-$, $-\text{C}\equiv\text{C}-\text{CH}_2-$ or $-\text{CH}_2-\text{C}\equiv\text{C}-$;

Z is



wherein R₄ and R₅ are hydrogen, hydroxy, halogen, lower alkyl, lower alkoxy or hydroxy(lower)alkyl, wherein R₄ and R₅ are not hydroxy and lower alkoxy at the same time;

R₁ is a saturated or unsaturated bivalent lower or medium aliphatic hydrocarbon residue, which is unsubstituted or substituted with halogen, alkyl, hydroxy, oxo, aryl selected from the group consisting of phenyl, tolyl and xylyl which is unsubstituted or substituted or heterocyclic group selected from the group consisting of furyl, thienyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, imidazolyl, pyrazolyl, furazanyl, pyranal, pyridyl, pyridazinyl, pyrimidyl, pyrazolyl, 2-pyrrolinyl, pyrrolidinyl, 2-imidazolinyl, imidazolidinyl, 2-pyrazolinyl, pyrazolidinyl, piperidino, piperazinyl, morpholino, indolyl, benzothienyl, quinolyl, isoquinolyl, purinyl, quinazolinyl, carbazolyl, acridinyl, phenanthridinyl, benzimidazolyl, benzimidazolinyl,

benzothiazolyl and phenothiazinyl which is unsubstituted or substituted, and at least one of carbon atom in the aliphatic hydrocarbon is optionally substituted by oxygen, nitrogen or sulfur; and

Ra is a saturated or unsaturated lower or medium aliphatic hydrocarbon residue which is unsubstituted or substituted with and may have a further substituent selected from the group consisting of oxo, hydroxy, lower alkoxy, lower alkanoyloxy, cyclo(lower)alkyl, cyclo(lower)alkyloxy, aryl, aryloxy, heterocyclic group and heterocyclic-oxy group; lower alkoxy; lower alkanoyloxy; cyclo(lower)alkyl; cyclo(lower)alkyloxy; aryl; aryloxy; heterocyclic group; heterocyclic-oxy.

19. (new): The method as described in Claim 1, wherein said prostaglandin compound is a 15-keto-16-mono or di-halogen prostaglandin compound.